

A regional checklist of Alysiinae (Hymenoptera: Braconidae) from Iran

Nahid Khajeh¹, Zahra Yari¹, Ehsan Rakhshani^{1*} and Francisco Javier Peris-Felipo²

1. Department of Plant Protection, College of Agriculture, University of Zabol, Zabol, P. O. Box: 98615-538, I. R. Iran.

2. Laboratory of Entomology and Pest Control, Institute Cavanilles of Biodiversity and Evolutionary Biology, University of Valencia, c/. Catedrático José Beltrán n 2, 46980 Paterna, Valencia, Spain.

Abstract: The present paper provides the updated checklist of the genera and species in the subfamily Alysiinae (Hymenoptera: Braconidae) from Iran. A total of 55 species belonging to 13 genera have been listed, that had been recorded from 15 provinces. The recorded species were belong to the genera *Alloea* Haliday, 1833 (one species), *Aphaereta* Foerster, 1862 (two species), *Aspilota* Foerster, 1862 (two species), *Dinotrema* Foerster, 1862 (four species), *Idiasta* Foerster, 1862 (one species), *Orthostigma* Ratzeburg, 1844 (three species), *Phaenocarpa* Foerster, 1861 (one species), *Pseudopezomachus* Montero, 1905 (one species) and *Synaldis* Foerster, 1862 (four species) from the tribe Alysiini, and *Chorebus* Haliday, 1833 (28 species), *Coelinidea* Viereck 1913 (one species), *Dacnusa* Haliday, 1833 (five species) and *Protodacnusa* Griffiths, 1964 (two species) from the tribe Dacnusiini.

Keywords: Alysiini, Dacnusiini, fauna, biological control, parasitoids.

Introduction

The subfamily Alysiinae is considered as a monophyletic group of parasitic braconids, on the basis of their broad exodont mandibles and the total loss of the occipital and epicnemal carinae (Wharton, 1997, 2002). All alysiines are koinobiont endoparasitoids of Cyclorrhaphous Diptera (Shaw and Huddleston, 1991). The adult female lays her eggs into the egg or larva of the host fly, and the progenies emerge from the host puparium. The outwardly directed and non-overlapping mandibles are using for emergence of the adults from the host puparium (Wharton, 1984). About 2,000 species and 104 genera have been recorded worldwide within Alysiinae, of which more than 65 genera and

1200 species occurred in the Palaearctic region (Yu *et al.*, 2012). This subfamily is traditionally divided in two large and polymorphic tribes Alysiini Leach, 1815 and Dacnusiini Foerster, 1862, which can be distinguished by the presence (Alysiini) or absence (Dacnusiini) of the vein r in the fore wings (Shenefelt, 1974; Docavo *et al.*, 2002).

From a biological point of view, members of the tribe Alysiini interact with a wide range of hosts from more than twenty families of Cyclorrhaphous Diptera, but the majority of the hosts belong to Anthomyiidae, Calliphoridae, Drosophilidae, Lonchopteridae, Muscidae, Phoridae, Sarcophagidae and Scathophagidae (Shenefelt, 1974; Belokobylskij and Kostromina, 2011; Yu *et al.*, 2012), often in humid habitats and ephemeral substrates (Wharton, 1984). On the contrary, member of the tribe Dacnusiini are almost exclusively specialized in leaf and stem miners from Agromyzidae, Chloropidae, Drosophilidae,

Handling Editor: Dr. Ali Asghar Talebi

* **Corresponding author**, e-mail: rakhshani@uoz.ac.ir

Received: 14 February 2014, Accepted: 4 March 2014

Published online: 5 March 2014

Ephydriidae and Pisilidae (Shaw and Huddleston, 1991; Belokobylskij and Kostromina, 2011; Peris-Felipo and Jiménez-Peydró, 2013).

Many species of the subfamily Alysiinae have been considered as biological control agents of the economically important Diptera pests (Griffiths, 1968; Drea *et al.*, 1986; Wharton, 1997; Achterberg *et al.*, 2012). A few attempts have also been done for control of the synanthropic fly populations (Calliphoridae, Muscidae and Sarcophagidae) by rearing and release of the genera *Alysia* Laterille, 1804, *Aphaereta* Foerster, 1862 and *Gnathopleura* Fischer, 1975 (Wharton, 1984, 1997; Marchiori, 2007; Henrique, 2006).

While the subfamily Alysiinae is a rather common group of the braconids in various habitats, they were greatly ignored in previous Braconidae lists (Fallahzadeh and Saghaei, 2010), until recent faunistic and taxonomic studies (Ghahari *et al.*, 2009; 2010a, b, 2011a, b, c, d; Ghahari and Fischer, 2011a, b; Fischer *et al.*, 2011; Lashkari-Bod *et al.*, 2011; Rastegar *et al.*, 2012). Within this context, the present work offers the updated checklist of Alysiinae of Iran with the main objective to provide a framework for the subsequent studies on this group, which lead to exploration and identification of many more species from other regions of Iran.

Materials and Methods

The checklist of previously recorded species was extracted from all available literatures (Ghahari *et al.*, 2009, 2010a, b, 2011a, b, c, d; Ghahari and Fischer, 2011a, b; Fischer *et al.*, 2011; Lashkari-Bod *et al.*, 2011; Rastegar *et al.*, 2012; Sedighi *et al.*, 2014). Some species were examined in the Naturhistorisches Museum Wien (Vienna, Austria; NHMW) and the British Natural History Museum (London, UK; BNHM). Classification, nomenclature and the distributional data of Braconidae followed Yu *et al.* (2012).

The species list of each tribe is sorted alphabetically based on the valid names. Data about the distribution in Iran and the general distribution in the world is also provided for each species, separately. A map is provided, indicating the distribution of Alysiinae species in different provinces of Iran (Fig. 1).

Results

In the current study, 55 species of Alysiinae belonging to 13 genera, including nine genera of the tribe Alysiini (*Alloea* Haliday, 1833, *Aphaereta* Foerster, 1862, *Aspilota* Foerster, 1862, *Idiasta* Foerster, 1862, *Dinotrema* Foerster, 1862, *Orthostigma* Ratzeburg, 1844, *Phaenocarpa* Foerster, 1861, *Pseudopezomachus* Montero, 1905 and *Synaldis* Foerster, 1862) and four genera of tribe Dacnusiini (*Chorebus* Haliday, 1833, *Coelinidea* Viereck 1913, *Dacnusa* Haliday, 1833 and *Protodacnusa* Griffiths, 1964) are listed. The species list of Alysiinae of Iran is provided below.

Subfamily Alysiinae Leach, 1815

Tribe Alysiini Leach, 1815

1. *Alloea contracta* (Haliday, 1833)

Distribution in Iran: Mazandaran province: Ghaemshahr, Savadkooh (Ghahari *et al.*, 2010a).

General distribution: Eastern Palaearctic and Western Palaearctic.

2. *Aphaereta difficilis* Nixon, 1939

Distribution in Iran: Ilam province: Moosiyan (Ghahari *et al.*, 2011b).

General distribution: Eastern Palaearctic and Western Palaearctic.

3. *Aphaereta minuta* (Nees, 1811)

Material examined: Iran: Tehran province, viii.1989, 54♀ and 11♂, ex pupa of Diptera ex snails, sp.1, CIE A20881, leg.: A. Nasrollahi (BNHM).

Distribution in Iran: Isfahan province: Aran-Bidgol and Semirom (Ghahari *et al.*, 2011a), Tehran province: Tehran.

General distribution: Eastern Palaearctic and Western Palaearctic.

Remark: This species has previously been recorded from Iran as *Asobara minuta* (Nees, 1811) (Ghahari *et al.*, 2011a).

4. *Aspilota alfalfae* Fischer, Lashkari-Bod, Rakhshani & Talebi, 2011

Material examined: Iran: Fars province: Zarghan, 20.v.2008, 1♀, swept on *Medicago sativa* L., leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Fars province: Zarghan (Fischer *et al.*, 2011).

General distribution: Iran.

5. *Aspilota delicata* Fischer, 1973

Distribution in Iran: Ardabil province: Dashte Moghan (Ghahari *et al.*, 2011c).

General distribution: Western Palaearctic.

6. *Dinotrema concinnum* (Haliday, 1838)

Distribution in Iran: Qazvin province: Danesfahan (Ghahari *et al.*, 2011d).

General distribution: Eastern Palaearctic and Western Palaearctic.

7. *Dinotrema cratocerum* (Thomson, 1895)

Distribution in Iran: Golestan province: Minoodasht (Ghahari *et al.*, 2010b).

General distribution: Eastern Palaearctic and Western Palaearctic.

8. *Dinotrema intermissum* (Fischer, 1974)

Distribution in Iran: Guilan province: Astara (Ghahari and Fischer, 2011a).

General distribution: Western Palaearctic.

9. *Dinotrema significarium* (Fischer, 1973)

Distribution in Iran: Sistan & Baluchestan province: Zabol (Ghahari *et al.*, 2010a).

General distribution: Eastern Palaearctic and Western Palaearctic.

10. *Idiasta picticornis* (Ruthe 1854)

Distribution in Iran: Sistan & Baluchestan province (Sedighi *et al.*, 2014).

General distribution: Eastern Palaearctic and Western Palaearctic.

11. *Orthostigma beyarslani* Fischer, 1995

Material examined: Iran: Fars province: Marvdasht, 09.v.2008, 2♀, swept on *Medicago sativa*, leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Fars province: Marvdasht (Fischer *et al.*, 2011; Lashkari-Bod *et al.*, 2011).

General distribution: Western Palaearctic.

12. *Orthostigma laticeps* (Thomson, 1895)

Distribution in Iran: Isfahan province: Isfahan (Ghahari *et al.*, 2011a)

General distribution: Eastern Palaearctic and Western Palaearctic.

13. *Orthostigma maculipes* (Haliday, 1838)

Distribution in Iran: Isfahan province: Isfahan (Ghahari *et al.*, 2011a)

General distribution: Eastern Palaearctic and Western Palaearctic.

14. *Phaenocarpa ruficeps* (Nees, 1812)

Distribution in Iran: Khorasan Razavi province: Mashhad (Ghahari *et al.*, 2011c)

General distribution: Eastern Palaearctic, Nearctic, Oriental and Western Palaearctic.

15. *Pseudopezomachus masii* Nixon, 1940

Distribution in Iran: Ilam province: Ilam (Ghahari *et al.*, 2011b).

General distribution: Western Palaearctic.

16. *Synaldis concolor* (Nees, 1812)

Material examined: Iran: Fars province: Neyriz, 13.iii.2009, 1♀, swept on *Triticum aestivum* L., leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Fars province: Neyriz (Lashkari-Bod *et al.*, 2011), Qazvin province: Danesfahan (Ghahari *et al.*, 2011d), East Azarbaijan province: Oskoo (Rastegar *et al.*, 2012).

General distribution: Eastern Palaearctic and Western Palaearctic.

17. *Synaldis distracta* (Nees, 1834)

Distribution in Iran: Isfahan province: Isfahan and Shahreza (Ghahari *et al.*, 2011a).

General distribution: Eastern Palaearctic, Oriental and Western Palaearctic.

18. *Synaldis maxima* (Fischer, 1962)

Distribution in Iran: Ilam province: Darrehshahr (Ghahari *et al.*, 2011b).

General distribution: Western Palaearctic.

19. *Synaldis megastigma* Fischer, 1967

Distribution in Iran: Guilan province: Minoodasht (Ghahari *et al.*, 2011c).

General distribution: Western Palaearctic.

Tribe Dacnusiini Foerster, 1862**20. *Chorebus (Chorebus) affinis*** (Nees, 1812)

Material examined: Iran: Fars province: Maharlu, 31.iii.2008, 1♀, weed, leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Mazandaran province: Chalous (Ghahari *et al.*, 2010b), Fars province: Maharlou (Lashkari-Bod *et al.*, 2011), Khorasan Razavi province: Kashmar (Ghahari *et al.*, 2011c).

General distribution: Eastern Palaearctic and Western Palaearctic.

Remark: This species has previously been recorded from Iran as *Chorebus longicornis* (Nees, 1811) (Ghahari *et al.*, 2010b).

21. *Chorebus (Stiphrocera) asphodeli* Griffiths, 1968

Distribution in Iran: Qazvin province: Moallem Kelāyeh (Ghahari *et al.*, 2011d), Ardabil province: Ardabil (Rastegar *et al.*, 2012).

General distribution: Western Palaearctic.

22. *Chorebus (Chorebus) axillaris* Fischer, Lashkari-Bod, Rakhshani & Talebi, 2011

Material examined: Iran: Fars province: Bidzard, 30.iv.2006, 1♀, orchard, leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Fars province: Bidzard (Fischer *et al.*, 2011).

General distribution: Iran.

23. *Chorebus (Phaenolexis) bathyzonus* (Marshall, 1895)

Distribution in Iran: Sistan & Baluchestan province (Sedighi *et al.*, 2014).

General distribution: Eastern Palaearctic and Western Palaearctic.

24. *Chorebus (Phaenolexis) compressiventris* (Telenga, 1935)

Distribution in Iran: Ardabil province: Pars Abad and Germe (Ghahari and Fischer, 2011b).

General distribution: Western Palaearctic.

25. *Chorebus (Stiphrocera) cubocephalus* (Telenga, 1934)

Distribution in Iran: Sistan & Baluchestan province (Sedighi *et al.*, 2014).

General distribution: Eastern Palaearctic and Western Palaearctic.

26. *Chorebus (Stiphrocera) diremtus* (Nees, 1834)

Distribution in Iran: East Azarbaijan province: Arasbaran (Ghahari and Fischer, 2011b).

General distribution: Eastern Palaearctic and Western Palaearctic.

27. *Chorebus (Phaenolexis) femoratus* Tobias, 1962

Distribution in Iran: Ilam province: Ilam (Ghahari *et al.*, 2011b).

General distribution: Western Palaearctic.

28. *Chorebus (Stiphrocera) flavipes* (Goureau, 1851)

Distribution in Iran: Guilan province: Astara (Ghahari and Fischer, 2011a), Ardabil province: Dashte Moghan (Ghahari *et al.*, 2011c).

General distribution: Eastern Palaearctic and Western Palaearctic.

29. *Chorebus (Phaenolexis) fuscipennis* (Nixon, 1937)

Distribution in Iran: Isfahan province: Isfahan (Ghahari *et al.*, 2011a).

General distribution: Eastern Palaearctic and Western Palaearctic.

30. *Chorebus (Chorebus) gracilipes* (Thomson, 1895)

Distribution in Iran: Kordestan province: Sanandaj (Ghahari *et al.*, 2010a)

General distribution: Eastern Palaearctic and Western Palaearctic.

31. *Chorebus (Stiphocera) groschkei* Griffiths, 1967

Material examined: Iran: Fars province: Shiraz, 01.v.2007, 1♂, weed, leg.: A. Lashkari-Bod (NHMW); Fars province: Takhte-Rostam, 03.vi.2008, 1♂, weed, leg.: A. Lashkari-Bod (NHMW); Fars province: Neyriz, 13.iii.2009, 1♂, *Triticum aestivum*, leg.: A. Lashkari-Bod (NHMW); Fars province: Bidzard, 30.iv.2006, 1♂, orchard, leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Fars province (Lashkari-Bod *et al.*, 2011).

General distribution: Western Palaearctic.

32. *Chorebus (Phaenolexis) iridis* Griffiths, 1968

Distribution in Iran: Qazvin province: Qazvin (Ghahari *et al.*, 2011d).

General distribution: Western Palaearctic.

33. *Chorebus (Stiphocera) lar* (Morley, 1924)

Distribution in Iran: Isfahan province: Anarak (Ghahari *et al.*, 2011a), Sistan & Baluchestan province (Sedighi *et al.*, 2014).

General distribution: Eastern Palaearctic and Western Palaearctic.

34. *Chorebus (Phaenolexis) leptogaster* (Haliday, 1839)

Distribution in Iran: Golestan province: Azadshahr (Ghahari *et al.*, 2011c).

General distribution: Eastern Palaearctic and Western Palaearctic.

35. *Chorebus (Chorebus) longiarticulis* Fischer, Lashkari-Bod, Rakhshani & Talebi, 2011

Material examined: Iran: Fars province: Zarghan, 27.iv.2008, 1♀, swept on *Medicago sativa*, leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Fars province: Zarghan (Fischer *et al.*, 2011).

General distribution: Iran.

36. *Chorebus (Stiphocera) misellus* (Marshall, 1895)

Distribution in Iran: Semnan province: Shahrood (Ghahari *et al.*, 2010a).

General distribution: Western Palaearctic.

37. *Chorebus (Stiphocera) mucronatus* (Telenga, 1935)

Distribution in Iran: Mazandaran province: Ramsar (Ghahari *et al.*, 2010a), Ilam province: Arkvaze Malekshahi (Ghahari *et al.*, 2011b).

General distribution: Eastern Palaearctic and Western Palaearctic.

38. *Chorebus (Chorebus) nigridiremptus* Fischer, Lashkari-Bod, Rakhshani & Talebi, 2011

Material examined: Iran: Fars province: Zarghan, 20.v.2008, 1♀, swept on *Medicago sativa*, leg.: A. Lashkari-Bod (NHMW); same locality, 22.vi.2008, 1♀, leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Fars province: Zarghan (Lashkari-Bod *et al.*, 2011).

General distribution: Iran.

39. *Chorebus (Chorebus) nixonii* Burgehele, 1959

Distribution in Iran: East Azarbaijan province: Jolfa (Ghahari and Fischer, 2011b).

General distribution: Western Palaearctic.

40. *Chorebus (Phaenolexis) ornatus* (Telenga, 1935)

Distribution in Iran: Qazvin province: Khoram Dasht (Ghahari *et al.*, 2011d).

General distribution: Western Palaearctic.

41. *Chorebus (Phaenolexis) posticus* Haliday, 1839

Distribution in Iran: Hamadan province: Hamadan (Ghahari *et al.*, 2009).

General distribution: Eastern Palaearctic, Nearctic and Western Palaearctic.

42. *Chorebus (Chorebus) properesam* Fischer, Lashkari-Bod, Rakhshani & Talebi, 2011

Material examined: Iran: Fars province: Zarghan, 14.vii.2008, 1♂, weed, leg.: A. Lashkari-Bod (NHMW), same locality, 09.v.2008, 1♂, swept on *Medicago sativa*, leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Fars province: Zarghan (Fischer *et al.*, 2011).

General distribution: Iran.

43. *Chorebus (Phaenolexis) stilifer* Griffiths, 1968

Material examined: Iran: Fars province: Marvdasht, 09.v.2008, 1♂, swept on *Medicago sativa*, leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Fars province: Marvdasht (Fischer *et al.*, 2011; Lashkari-Bod *et al.*, 2011).

General distribution: Western Palaearctic.

44. *Chorebus (Phaenolexis) tamsi* (Nixon, 1944)

Material examined: Iran: Fars province: Abadeh, 19.iv.2007, 1♀, weed, leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Fars province: Abadeh (Fischer *et al.*, 2011; Lashkari-Bod *et al.*, 2011).

General distribution: Western Palaearctic.

45. *Chorebus (Stiphrocera) uliginosus* (Haliday, 1839)

Distribution in Iran: Mazandaran province: Savadkooh (Ghahari *et al.*, 2010b).

General distribution: Eastern Palaearctic and Western Palaearctic.

46. *Chorebus (Stiphrocera) venustus* (Tobias, 1962)

Distribution in Iran: Isfahan province: Ardestan (Ghahari *et al.*, 2011a).

General distribution: Eastern Palaearctic and Western Palaearctic.

47. *Chorebus (Chorebus) zarghanensis* Fischer, Lashkari-Bod, Rakhshani & Talebi, 2011

Material examined: Iran: Fars province: Zarghan, 09.v.2008, 3♀ (including holotype), swept on *Medicago sativa*, leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Fars province: Zarghan (Fischer *et al.*, 2011).

General distribution: Iran.

48. *Coelinidea gracilis* (Curtis, 1829)

Material examined: Iran: Fars province: Abadeh, 19.iv.2007, 3♂, weed, leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Fars province: Abadeh (Fischer *et al.*, 2011, Lashkari-Bod *et al.*, 2011), Ilam province: Dehloran (Ghahari *et al.*, 2011b).

General distribution: Eastern Palaearctic and Western Palaearctic.

Remarks: This species has previously been recorded from Iran as *Lepton gracilis* (Curtis, 1829) (Fischer *et al.*, 2011, Lashkari-Bod *et al.*, 2011).

49. *Dacnusa (Pachysema) alpestris* Griffiths, 1967

Distribution in Iran: Ilam province: Abadan (Ghahari *et al.*, 2011b).

General distribution: Eastern Palaearctic and Western Palaearctic.

50. *Dacnusa (Dacnusa) confinis* Ruthe, 1859

Distribution in Iran: Qazvin province: Kouhin (Ghahari *et al.*, 2011d).

General distribution: Western Palaearctic.

51. *Dacnusa (Dacnusa) gentianae* Griffiths, 1967

Distribution in Iran: Isfahan province: Golpayegan and Isfahan (Ghahari *et al.*, 2011a).

General distribution: Western Palaearctic.

52. *Dacnusa (Aphanta) hospita* (Foerster, 1862)

Material examined: Iran: Fars province: Neyriz, 13.iii.2009, 1♀, swept on *Triticum aestivum*, leg.: A. Lashkari-Bod (NHMW).

Distribution in Iran: Fars province: Neyriz (Fischer *et al.*, 2011, Lashkari-Bod *et al.*, 2011).

General distribution: Eastern Palaearctic and Western Palaearctic.

53. *Dacnusa (Pachysema) sibirica* Telenga, 1935

Distribution in Iran: Ardabil province: Ardabil (Fathi, 2011), Sistan & Baluchestan province (Sedighi et al., 2014).

General distribution: Eastern Palaearctic, Oriental and Western Palaearctic.

54. *Protodacnusa aridula* (Thomson, 1895)

Distribution in Iran: Fars province: Shiraz (Ghahari et al., 2010a).

General distribution: Eastern Palaearctic and Western Palaearctic.

55. *Protodacnusa litoralis* Griffiths, 1964

Distribution in Iran: Qazvin province: Takestan (Ghahari et al., 2011d).

General distribution: Eastern Palaearctic and Western Palaearctic.



- | | | | |
|-----------------------------------|--------------------------------------|------------------------------------|-----------------------------------|
| 1. <i>Alloea contracta</i> | 16. <i>Synaldis concolor</i> | 31. <i>Chorebus groschkei</i> | 46. <i>Chorebus venustus</i> |
| 2. <i>Aphaereta difficilis</i> | 17. <i>Synaldis distracta</i> | 32. <i>Chorebus iridis</i> | 47. <i>Chorebus zarghanensis</i> |
| 3. <i>Aphaereta minuta</i> | 18. <i>Synaldis maxima</i> | 33. <i>Chorebus lar</i> | 48. <i>Coelinidea gracilis</i> |
| 4. <i>Aspilota alfalvae</i> | 19. <i>Synaldis megastigma</i> | 34. <i>Chorebus leptogaster</i> | 49. <i>Dacnusa alpestris</i> |
| 5. <i>Aspilota delicata</i> | 20. <i>Chorebus affinis</i> | 35. <i>Chorebus longiarticulis</i> | 50. <i>Dacnusa confinis</i> |
| 6. <i>Dinotrema concinnum</i> | 21. <i>Chorebus asphodeli</i> | 36. <i>Chorebus misellus</i> | 51. <i>Dacnusa gentianae</i> |
| 7. <i>Dinotrema cratocerum</i> | 22. <i>Chorebus axillaris</i> | 37. <i>Chorebus mucronatus</i> | 52. <i>Dacnusa hospita</i> |
| 8. <i>Dinotrema intermissum</i> | 23. <i>Chorebus bathyzonus</i> | 38. <i>Chorebus nigridiremptus</i> | 53. <i>Dacnusa sibirica</i> |
| 9. <i>Dinotrema significarium</i> | 24. <i>Chorebus compressiventris</i> | 39. <i>Chorebus nixonii</i> | 54. <i>Protodacnusa aridula</i> |
| 10. <i>Idiasta picticornis</i> | 25. <i>Chorebus cubocephalus</i> | 40. <i>Chorebus ornatus</i> | 55. <i>Protodacnusa litoralis</i> |
| 11. <i>Orthostigma beyarslani</i> | 26. <i>Chorebus diremtus</i> | 41. <i>Chorebus posticus</i> | |
| 12. <i>Orthostigma laticeps</i> | 27. <i>Chorebus femoratus</i> | 42. <i>Chorebus properesam</i> | |
| 13. <i>Orthostigma maculipes</i> | 28. <i>Chorebus flavipes</i> | 43. <i>Chorebus stilifer</i> | |
| 14. <i>Phaenocarpa ruficeps</i> | 29. <i>Chorebus fuscipennis</i> | 44. <i>Chorebus tamsi</i> | |
| 15. <i>Pseudopezomachus masii</i> | 30. <i>Chorebus gracilipes</i> | 45. <i>Chorebus uliginosus</i> | |

Figure 1 Distribution map for Alysiniinae species in different provinces of Iran. The numbers on the map correspond to the species names which are listed below.

Discussion

Considering the vast number of the known Alysiinae species in Palaearctic region (Yu *et al.*, 2012), occurrence of only 55 species is clearly indicating the poor knowledge about fauna of this group in Iran. Discrete and limited number of faunistic studies, mainly on the local sense (Ghahari *et al.*, 2009, 2010, 2011, 2012a, b; Fischer *et al.*, 2011 Lashkari-Bod *et al.*, 2011; Rastegar *et al.*, 2012) have yet to be expanded with explorations of many more areas in other parts of the country. An overall comparison of the fauna of Iranian Alysiinae with the neighboring countries (Tobias, 1986; Fischer and Beyarslan, 2012) is also indicating the existence of many other genera and species to be detected.

While, number of the recorded genera was greater in the tribe Alysiini (9 genera and 19 species), more species were recorded from the tribe Dacnusi (4 genera and 36 species). Among the genera of the later group, the genus *Chorebus* includes the majority of taxa (28 species). This genus with approximately 215 Holarctic species, is the largest genus of the Dacnusi (Tobias, 1986).

Except for a few endemic species (Fischer *et al.*, 2011) it seems that the rest species are widely distributed in the Palaearctic region. A few species as *Phaenocarpa ruficeps* and *Synaldis distracta* have also been recorded from the Oriental region. There are some records of *Phaenocarpa ruficeps* and *Chorebus posticus* from the Nearctic region. Meanwhile, the occurrence of some species, including *Dinotrema intermissum* (Austria, Iran, Turkey), *Orthostigma beyarslani* (Iran, Spain, Turkey), *Pseudopezomachus masii* (Greece, Iran, Libya) and *Chorebus stilifer* (Germany, Iran) is documented in a very few countries in the western Palaearctic region (Yu *et al.*, 2012). The fragmented type of the recorded localities, suggests a wider distribution of these species in the Western Palaearctic region, including the central Asian area.

In the same way, very little information is available about host association of the recorded species in Iran. *Dacnusa sibirica* was reared from the leafminer, *Chromatomyia horticola* (Goureau) (Diptera: Agromyzidae), one of the most important pest of canola, *Brassica napus* L., crop in Ardabil region (Fathi, 2011). This species is a commercially mass reared biological control agents of the agromyzid leafminers (Lenteren, 2003; Abd-Rabou, 2006).

In General, the limited number of the known Alysiinae from Iran reflecting the paucity of the researches on these group, as well as their relative rarity. Further investigations both on the fauna and host association of the Iranian Alysiinae are necessary to provide the basis for biological control of the dipterous pests in agricultural and urban landscapes.

Acknowledgements

This work has supported by the grant No. 89-9198, University of Zabol. We would like to thank Dominique Zimmermann and Manuela Vizek from Naturhistorisches Museum of Wien for their kindness and help with the loan of some specimens.

References

- Abd-Rabou, S. 2006. Biological control of the leafminer, *Liriomyza trifolii* by introduction, releasing, evaluation of the parasitoids *Diglyphus isaea* and *Dacnusa sibirica* on vegetables crops in greenhouses in Egypt. Archives of Phytopathology and Plant Protection, 39 (6): 439-443.
- Achterberg, C. van, Teixeira, T. and Oliveira, L. 2012. *Aphaereta ceratitivora* sp. n. (Hymenoptera, Braconidae), a new parasitoid of *Ceratitis capitata* (Wiedemann) (Diptera, Tephritidae) from the Azores. Zookeys, 222: 1-9.
- Belokobylskij, S. A. and Kostromina, T. S. 2011. Two braconid alysiine genera new for Russia. Zoosystematica Rossica, 20 (1): 85-95.

- Docavo, I., Tormos, J. and Fischer, M. 2002. Three new species of *Chorebus* from Spain (Hymenoptera: Braconidae: Alysiinae). *Florida Entomologist*, 85 (1): 208-215.
- Drea, J. J. and Hendrickson, R. M. 1986. Analysis of a successful classical biological control project: the alfalfa blotch leafminer (Diptera: Agromyzidae) in the northeastern United States. *Environmental Entomology*, 15 (3): 448-455.
- Fallahzadeh, M. and Saghaei, N. 2010. Checklist of Braconidae (Insecta: Hymenoptera) from Iran. *Munis Entomology & Zoology*, 5: 170-186.
- Fathi, S. A. A. 2011. Tritrophic interactions of nineteen canola cultivars - *Chromatomya horticola*-parasitoids in Ardabil region. *Munis Entomology & Zoology*, 6: 449-454.
- Fischer, M. and Beyarslan, A. 2012. New species of *Synaldis* Foerster and *Idiasta* Foerster, and further records of Turkish Alysiini: (Hymenoptera: Braconidae: Alysiinae). *Zoology in the Middle East*, 55 (1): 55-64.
- Fischer, M., Lashkari-Bod, A., Rakhshani, E. and Talebi, A. A. 2011. Alysiinae from Iran (Insecta: Hymenoptera: Braconidae: Alysiinae). *Annalen des Naturhistorischen Museums in Wien, serie B*, 112: 115-132.
- Ghahari, H. and Fischer, M. 2011a. A study on the Braconidae (Hymenoptera: Ichneumonoidea) from some regions of northern Iran. *Entomofauna*, 32 (8): 181-196.
- Ghahari, H. and Fischer, M. 2011b. A contribution to the Braconidae (Hymenoptera: Ichneumonoidea) from north-western Iran. *Calodema*, 134: 1-6.
- Ghahari, H., Fischer, M., Erdogan, O. Ç., Beyarslan, A., Hedqvist, K. J. and Ostovan, H. 2009. Faunistic note on the Braconidae (Hymenoptera: Ichneumonoidea) in Iranian alfalfa fields and surrounding grasslands. *Entomofauna*, 30 (24): 437-444.
- Ghahari, H., Fischer, M., Hedqvist, K. J., Erdogan, O. Ç., Achterberg, C. van and Beyarslan, A. 2010a. Some new records of Braconidae (Hymenoptera) for Iran. *Linzer Biologische Beiträge*, 42 (2): 1395-1404.
- Ghahari, H., Fischer, M., Endogan, O. Ç., Beyarslan, A. and Ostovan, H. 2010b. A contribution to the braconid wasps (Hymenoptera: Braconidae) from the forests of northern Iran. *Linzer Biologische Beiträge*, 42 (1): 621-634.
- Ghahari, H., Fischer, M. and Papp, J. 2011a. A study on the braconid wasps (Hymenoptera: Braconidae) from Isfahan province, Iran. *Entomofauna*, 32 (16): 261-272.
- Ghahari, H., Fischer, M. and Papp, J. 2011b. A study on the Braconidae (Hymenoptera: Ichneumonoidea) from Ilam province, Iran. *Calodema*, 160: 1-5.
- Ghahari, H., Fischer, M., Sakenin, H. and Imani, S. 2011c. A contribution to Agathidinae, Alysiinae, Aphidiinae, Braconinae, Microgastrinae and Opiinae (Hymenoptera, Braconidae) from cotton fields and surrounding grassland of Iran. *Linzer Biologische Beiträge*, 43 (2): 1269-1276.
- Ghahari, H., Fischer, M. and Papp, J. 2011d. A study on the Braconidae (Hymenoptera, Ichneumonoidea) from Qazvin province, Iran. *Entomofauna*, 32 (9): 197-204.
- Griffiths, G. C. D. 1968. The Alysiinae (Hym., Braconidae) parasites of the Agromyzidae (Diptera). *Beitraege zur Entomologie*, 18:5-62.
- Henrique, C. 2006. *Gnathopleura quadridentata* (Wharton) (Hymenoptera: Braconidae: Alysiinae) as natural enemy of *Sarcodexia Lambens* (Wiedemann) (Diptera: Sarcophagidae) in Brazil. *Brazilian Journal of Veterinary Research and Animal Science*, 43 (5): 708-710.
- Lashkari-Bod, A., Rakhshani, E., Talebi, A. A., Lozan, A. and Zikic, V. 2011. Contribution to the knowledge of Braconidae (Hym., Ichneumonoidea) of Iran. *Biharean Biologist*, 5 (2): 147-150.
- Lenteren, J. C. van. 2003. Commercial availability of biological control agents. In: Lenteren, J. C. van (Ed.), *Quality Control and Production of Biological Control Agents: Theory and Testing Procedures*, CAB International, Wallingford, UK, pp: 167-179.
- Marchiori, C. H. 2007. *Gnathopleura quadridentata* Wharton, 1986 (Hymenoptera;

- Braconidae; Alysiinae) and their hosts collected in different substrates in Caldas Novas, Goiás. *Brazilian Journal of Biology*, 67 (1): 101-103.
- Peris-Felipo F. J. and Jiménez-Peydró R. 2013. Contribution to the phenological knowledge of *Aspilota*-group (Hymenoptera Braconidae Alysiinae) in Mediterranean landscapes. *Biodiversity Journal*, 4 (3): 419-426.
- Rastegar, J., Sakenin, H., Khodaparast, S. and Havaskary, M. 2012. On a collection of Braconidae (Hymenoptera) from East Azarbaijan and vicinity, Iran. *Calodema*, 226: 1-4.
- Sedighi, S., Madjdzadeh, S. M. and Rakhshani, E. 2014. A survey on Alysiinae (Hym.: Braconidae) associated with Alfalfa in central part of Sistan Baluchistan Province. Third Integrated Pest Management Conference (IPMC), 21–22 January 2014, Kerman, Iran.
- Sharkey, M. J. and Wharton, R. A. 1997. Morphology and terminology. In: Wharton, R. A., Marsh, P. M. and Sharkey, M. J. (Eds), *Manual of the New World Genera of the Family Braconidae (Hymenoptera)*. International Society Hymenopterists, Washington, D. C. pp: 19-37.
- Shaw, M. R. and Huddleston, T. 1991. Classification and biology of braconid wasps (Hymenoptera: Braconidae). *Handbooks for the Identification of British Insects*, 7 (11), 1-26.
- Shenefelt, R. D. 1974. Braconidae 7. Alysiinae. In: Van der Vecht, J. and Shenefelt, R. D. (Eds.), *Hymenopterorum Catalogus, Part II*. Dr. W. Junk: The Hague. pp: 937-1113.
- Tobias, V. I. 1986. Subfamily Alysiinae. In: Medvedev G.S. (Ed.), *Keys to the Insects of the European Part of the USSR, III. Part V*. Nauka Publisher, Leningrad. (in Russian, English translation in 1995). pp: 156-386.
- Wharton, R. A. 1984. Biology of the Alysiini (Hymenoptera: Braconidae), parasitoids of cyclorrhaphous Diptera. *Texas Agricultural Experiment Station Technical Monograph* 11, 1-39.
- Wharton, R. A. 1997. Subfamily Alysiinae. In: Wharton, R. A., Marsh, P. M. and Sharkey, M. J. (Eds.), *Manual of the New World Genera of the Family Braconidae (Hymenoptera)*. International Society Hymenopterists, Washington, D. C. pp: 84-116.
- Wharton, R. A. 2002. Revision of the Australian Alysiini (Hymenoptera: Braconidae). *Invertebrate Systematics*, 16 (1): 7-105.
- Yu, D. S., Achterberg, C. van and Horstmann, K. 2012. *World Ichneumonoidea 2011. Taxonomy, biology, Morphology and distribution (Braconidae)*. Taxapad (Scientific names for information management) Interactive catalogue on DVD/CDROM. Vancouver.

چک لیست زنبورهای زیرخانواده Alysiniinae (Hymenoptera: Braconidae) ایران

ناهید خواجه^۱، زهرا یاری^۱، احسان رخشانی^{۱*} و فرانسیسکو ژاوییر پریس فلیپو^۲

۱- گروه گیاه پزشکی، دانشکده کشاورزی، دانشگاه زابل، زابل، ایران.

۲- مرکز مطالعات بیولوژی فرگشتی و تنوع گونه‌ای کوانیلِس، دانشگاه والنسیا، اسپانیا.

* پست الکترونیکی نویسنده مسئول مکاتبه: rakhshani@uoz.ac.ir

دریافت: ۲۵ بهمن ۱۳۹۲؛ پذیرش: ۱۳ اسفند ۱۳۹۲

چکیده: در این مقاله فهرست جنس‌ها و گونه‌های زیرخانواده Alysiniinae (Hymenoptera: Braconidae) ایران ارائه شده است. در مجموع تعداد ۵۵ گونه متعلق به ۱۳ جنس از ۱۵ استان کشور فهرست شدند. گونه‌های ثبت شده متعلق به جنس‌های *Alloea* Haliday, 1833 (یک گونه)، *Aphaereta* Foerster, 1862 (دو گونه)، *Aspilota* Foerster, 1862 (دو گونه)، *Dinotrema* Foerster, 1862 (چهار گونه)، *Idiasta* Foerster, 1862 (یک گونه)، *Orthostigma* Ratzeburg, 1844 (سه گونه)، *Phaenocarpa* Foerster, 1861 (یک گونه)، *Pseudopezomachus* Montero, 1905 (یک گونه) و *Synaldis* Foerster, 1862 (چهار گونه) از قبیله Alysiniinae و جنس‌های *Chorebus* Haliday, 1833 (۲۸ گونه)، *Coelinidea* Viereck 1913 (یک گونه)، *Dacnusa* Haliday, 1833 (پنج گونه) و *Protodacnusa* Griffiths, 1964 (دو گونه) از قبیله Dacnusiinae بودند.

واژگان کلیدی: Alysiniinae، Dacnusiinae، فون، کنترل بیولوژیک، پارازیتوئید.